

## GUIDELINES FOR PREPARATION OF ABSTRACTS

### GUIDELINES

To be selected for either an oral or poster presentation authors must submit an abstract.

- Abstracts should be prepared in Word using Calibri 12-point font.
- Select A4 page size with a 2.5 cm margin all around.
- The title should be typed in BOLD CAPITALS, 14-point font, centered, leaving a gap of one line after.
- Authors' names should be in lower case, centered, leaving a gap of one line after.
- Underline the name of the presenting author.
- Authors' affiliations and addresses should appear after a one-line space in lower case, aligned to the left, with multiple affiliations indicated by superscripted numbers.
- Leave a double space and proceed with the abstract, using single spacing and the writing justified without indentation.

### SAMPLE ABSTRACT FORMAT

#### HMW GLUTENIN SUBUNITS FROM 1S<sup>1</sup> GENOME OF *AEGILOPS LONGISSIMA*

Wujun Ma<sup>1</sup>, Yueming Yan<sup>2</sup>, Shunli Wang<sup>1,2</sup>, Zitong Yu<sup>1,2</sup>, H Weißgerber<sup>3</sup>, FJ Zeller<sup>3</sup>, SLK Hsam<sup>4</sup>,  
Dean Diepeveen<sup>1</sup>, Shahidul Islam<sup>1</sup>, Rudi Appels<sup>1</sup>

<sup>1</sup>Australian Export Grain Innovation Centre, <sup>2</sup>Baron-Hay Court, Perth, WA 6151, & School of Veterinary and Life Sciences, Murdoch University, South St, Perth WA 6150

<sup>3</sup>College of Life Science, Capital Normal University, 100048 Beijing, China

<sup>4</sup>Division of Plant Breeding and Applied Genetics, Technical University of Munich, D-85350 Freising-Weihenstephan, Germany

#### ABSTRACT

A Chinese Spring (CS) substitution line was produced in which a chromosome (1S<sup>1</sup>) from *Aegilops longissima* substituted one chromosome (1B) of CS with a significantly increase of dough rheological property. A range of studies have been carried out in aspects of high molecular weight glutenin coding genes, protein accumulations, protein bodies, starch granules and protein disulfide isomerase (PDI) and PDI-like protein expressions (PDI and PDI-like protein can control diversified metabolic functions) in order to uncover the mechanism underlying the superior dough and breadmaking quality of this substitution line.